

## Lecture 7: Spectral Expansion I

September 28, 2004

Scribe: Arash Rafiey

## 1 Spectral Expansion

Let  $G$  be a  $d$ -regular multigraph on  $n$  vertices and  $A$  be the normalized adjacency matrix of  $G$ . Set  $\lambda_2(G) = \max\{|\lambda_i| \mid i \geq 2\}$  where  $1 = \lambda_1 \geq \lambda_2 \geq \dots \geq \lambda_n$  are eigenvalues of  $G$ .

**Definition 1**  $G$  has spectral expansion  $\lambda$ , if  $\lambda_2(G) \leq \lambda$ .

**Theorem 2** Spectral expansion  $\Rightarrow$  Vertex expansion .

If  $G$  has spectral expansion  $\lambda$  then  $\forall \alpha < 1$ ,  $G$  has vertex expansion  $(\alpha n, \frac{1}{(1-\alpha)\lambda^2 + \alpha})$ .

We require some definitions and lemmas to prove the Theorem.

**Definition 3** For any probability distribution  $\pi$ , the collision probability of  $\pi$  is defined as:

$$Col(\pi) = \sum_{i=1}^{i=n} \pi_i^2.$$

**Definition 4** For any probability distribution  $\pi$ ,  $Supp(\pi) = \{i \mid \pi_i > 0\}$

In what follows let  $u = (\frac{1}{n}, \dots, \frac{1}{n})$ .  $u$  is uniform distribution.

**Lemma 5** For any distribution  $\pi$

$$1) Col(\pi) \geq \frac{1}{|Supp(\pi)|}.$$

$$2) Col(\pi) = \|\pi - u\|^2 + \frac{1}{n} = \|\pi - u\|^2 + Col(u).$$

**Remark 6** For distribution which is uniform over  $Supp(\pi)$ , the collision probability is

$$\sum_{i \in Supp(\pi)} \frac{1}{|Supp(\pi)|^2} = \frac{1}{|Supp(\pi)|}.$$

It means that among all probability distribution with same Support, the uniform distribution has minimum collision probability.

**Proof of Lemma 5:**

**Proof of part (1):** By Cauchy-Schwarz,  $\forall$  vectors  $u, v$ ,  $(u, v) \leq \|u\| \|v\|$ , where  $(u, v)$  is the inner product of  $u, v$ .

Define vector  $x$  as follow:

$$x_i = \begin{cases} 1 & \text{if } \pi_i > 0 \\ 0 & \text{if } \pi_i = 0 \end{cases} .$$

We have  $\sum x_i^2 = |Supp(\pi)|$ . Therefore  $1 = (\pi, x)^2 \leq \|\pi\|^2 \|x\|^2 = Col(\pi) |Supp(\pi)|$ . ■

**Proof of part (2):** Write  $\pi = u + \pi^\perp$ . Then  $Col(\pi) = \|\pi\|^2 = \|u + \pi^\perp\|^2 = (u + \pi^\perp, u + \pi^\perp) = (u, u) + 2(u, \pi^\perp) + (\pi^\perp, \pi^\perp) = \|u\|^2 + \|\pi^\perp\|^2 = \frac{1}{n} + \|\pi - u\|^2$ .

Note that  $(u, \pi^\perp) = 0$  from previous lecture! ■

### Proof of Theorem 2:

$\forall \pi, \|A\pi - u\|^2 \leq \lambda^2 \|\pi - u\|^2$ . Let  $\pi$  be uniform distribution on set  $S$  of size  $l \leq \alpha n$ . By Lemma 5  $|N(S)| = |Supp(A\pi)| \geq \frac{1}{Col(A\pi)}$ . So we need to upperbound  $Col(A\pi)$ .

$$Col(A\pi) = \|A\pi - u\|^2 + \frac{1}{n} \leq \lambda^2 \|\pi - u\|^2 + \frac{1}{n} = \lambda^2 (Col(\pi) - \frac{1}{n}) + \frac{1}{n}.$$

So we have

$$Col(A\pi) - \frac{1}{n} \leq \lambda^2 (Col(\pi) - \frac{1}{n}).$$

Finally,

$$\frac{1}{|N(S)|} - \frac{1}{n} \leq Col(A\pi) - \frac{1}{n} \leq \lambda^2 \left( \frac{1}{|S|} - \frac{1}{n} \right).$$

Therefore we have  $\frac{|S|}{|N(S)|} \leq \lambda^2 + |S| \frac{1-\lambda^2}{n}$ , and  $\frac{|N(S)|}{|S|} \geq \frac{n}{\lambda^2 n + |S|(1-\lambda^2)} \geq \frac{n}{\lambda^2 n + \alpha n(1-\lambda^2)} = \frac{1}{\lambda^2 + \alpha(1-\lambda^2)}$ . ■

### Theorem 7 Vertex expansion $\Rightarrow$ Spectral expansion [1]

$\forall$  constant  $\gamma > 0, d > 2$ , there is  $\lambda < 1$ , s.t. if  $G$  is a  $d$ -regular graph with  $n$  vertices and  $(\frac{n}{2}, 1 + \gamma)$ -expansion, then it has  $\lambda \leq 1 - \Omega(\frac{\gamma^2}{d^2})$  spectral expansion.

## References

- [1] N. Alon, Z. Galil and V. D. Milman, Better expanders and superconcentrators, J. Algorithms 8 (1987), 337-347.