

BAYESIAN NETWORKS

CHAPTER 14.1-3

Chapter 14.1-3 1

Outline

- ◇ Syntax
- ◇ Semantics
- ◇ Parameterized distributions

Bayesian networks

A simple, graphical notation for conditional independence assertions and hence for compact specification of full joint distributions

Syntax:


- a set of nodes, one per variable
- a directed, acyclic graph (link \approx "directly influences")
- a conditional distribution for each node given its parents:
 $P(X_i | Parents(X_i))$

In the simplest case, conditional distribution represented as a **conditional probability table (CPT)** giving the distribution over X_i for each combination of parent values

Chapter 14.1-3 2

Example

Topology of network encodes conditional independence assertions:



Weather is independent of the other variables

Toothache and *Catch* are conditionally independent given *Cavity*

Chapter 14.1-3 4

Example

I'm at work, neighbor John calls to say my alarm is ringing, but neighbor Mary doesn't call. Sometimes it's set off by minor earthquakes. Is there a burglar?


Variables: *Burglar*, *Earthquake*, *Alarm*, *JohnCalls*, *MaryCalls*

Network topology reflects "causal" knowledge:

- A burglar can set the alarm off
- An earthquake can set the alarm off
- The alarm can cause Mary to call
- The alarm can cause John to call

Chapter 14.1-3 5

Example contd.




Chapter 14.1-3 3

Chapter 14.1-3 6

Compactness

A CPT for Boolean X_i with k Boolean parents has 2^k rows for the combinations of parent values

Each row requires one number p for $X_i = true$ (the number for $X_i = false$ is just $1 - p$)



If each variable has no more than k parents, the complete network requires $O(n \cdot 2^k)$ numbers

I.e., grows linearly with n , vs. $O(2^n)$ for the full joint distribution


For burglary net, ?? numbers

Chapter 14.3 7

Global semantics

"Global" semantics defines the full joint distribution as the product of the local conditional distributions:

$$P(x_1, \dots, x_n) = \prod_{i=1}^n P(x_i | \text{parents}(X_i))$$



e.g., $P(j \wedge m \wedge a \wedge \neg b \wedge \neg e)$

$$= P(j|a)P(m|a)P(a|\neg b, \neg e)P(\neg b)P(\neg e)$$

$$= 0.9 \times 0.7 \times 0.001 \times 0.999 \times 0.998$$


$$\approx 0.000063$$

Chapter 14.3 10

Compactness

A CPT for Boolean X_i with k Boolean parents has 2^k rows for the combinations of parent values

Each row requires one number p for $X_i = true$ (the number for $X_i = false$ is just $1 - p$)



If each variable has no more than k parents, the complete network requires $O(n \cdot 2^k)$ numbers

I.e., grows linearly with n , vs. $O(2^n)$ for the full joint distribution

For burglary net, $1 + 1 + 4 + 2 + 2 = 10$ numbers (vs. $2^5 - 1 = 31$)

Chapter 14.3 8


Chapter 14.3 11

Global semantics

Global semantics defines the full joint distribution as the product of the local conditional distributions:

$$P(x_1, \dots, x_n) = \prod_{i=1}^n P(x_i | \text{parents}(X_i))$$

e.g., $P(j \wedge m \wedge a \wedge \neg b \wedge \neg e)$



Chapter 14.3 9

Constructing Bayesian networks

Need a method such that a series of locally testable assertions of conditional independence guarantees the required global semantics

1. Choose an ordering of variables X_1, \dots, X_n
2. For $i = 1$ to n
 - add X_i to the network
 - select parents from X_1, \dots, X_{i-1} such that $P(X_i | \text{Parents}(X_i)) = P(X_i | X_1, \dots, X_{i-1})$

This choice of parents guarantees the global semantics:

$$P(X_1, \dots, X_n) = \prod_{i=1}^n P(X_i | X_1, \dots, X_{i-1}) \quad (\text{chain rule})$$

$$= \prod_{i=1}^n P(X_i | \text{Parents}(X_i)) \quad (\text{by construction})$$

Chapter 14.3 12

Example

Suppose we choose the ordering M, J, A, B, E




$P(J|M) = P(J)$?

Chapter 11.1-3 13

Example

Suppose we choose the ordering M, J, A, B, E




$P(J|M) = P(J)$? No
 $P(A|J, M) = P(A|J)$? $P(A|J, M) = P(A)$? No
 $P(B|A, J, M) = P(B|A)$? Yes
 $P(B|A, J, M) = P(B)$? No
 $P(E|B, A, J, M) = P(E|A)$?
 $P(E|B, A, J, M) = P(E|A, B)$?

Chapter 11.1-3 16

Example

Suppose we choose the ordering M, J, A, B, E




$P(J|M) = P(J)$? No
 $P(A|J, M) = P(A|J)$? $P(A|J, M) = P(A)$?

Chapter 11.1-3 14

Example

Suppose we choose the ordering M, J, A, B, E




$P(J|M) = P(J)$? No
 $P(A|J, M) = P(A|J)$? $P(A|J, M) = P(A)$? No
 $P(B|A, J, M) = P(B|A)$? Yes
 $P(B|A, J, M) = P(B)$? No
 $P(E|B, A, J, M) = P(E|A)$? No
 $P(E|B, A, J, M) = P(E|A, B)$? Yes

Chapter 11.1-3 17


Example

Suppose we choose the ordering M, J, A, B, E



$P(J|M) = P(J)$? No
 $P(A|J, M) = P(A|J)$? $P(A|J, M) = P(A)$? No
 $P(B|A, J, M) = P(B|A)$?
 $P(B|A, J, M) = P(B)$?

Example contd.




Deciding conditional independence is hard in noncausal directions
 (Causal models and conditional independence seem hardwired for humans!)
 Assessing conditional probabilities is hard in noncausal directions
 Network is less compact: $1 + 2 + 4 + 2 + 4 = 13$ numbers needed

Chapter 11.1-3 15

Chapter 11.1-3 18

Example: Car diagnosis



Chapter 11.1-3 19

Compact conditional distributions contd.

Noisy-OR distributions model multiple noninteracting causes

- Parents $U_1 \dots U_k$ include all causes (can add leak node)
- Independent failure probability q_i for each cause alone


$$\Rightarrow P(X|U_1 \dots U_k, \neg U_{j+1} \dots \neg U_n) = 1 - \prod_{i=1}^k q_i$$

<i>Cold</i>	<i>Flu</i>	<i>Malaria</i>	$P(\text{Fever})$	$P(\neg \text{Fever})$
F	F	F	0.0	1.0
F	F	T	0.9	0.1
F	T	F	0.8	0.2
F	T	T	0.98	$0.02 = 0.2 \times 0.1$
F	T	F	0.4	0.6
T	F	T	0.94	$0.06 = 0.6 \times 0.1$
T	F	F	0.88	$0.12 = 0.6 \times 0.2$
T	T	T	0.988	$0.012 = 0.6 \times 0.2 \times 0.1$

Number of parameters **linear** in number of parents

Chapter 11.1-3 22


Example: Car insurance



Chapter 11.1-3 20

Hybrid (discrete+continuous) networks

Discrete (*Subsidy?* and *Buy?*): continuous (*Harvest* and *Cost*)



Option 1: discretization—possibly large errors, large CPTs
 Option 2: finitely parameterized canonical families

- Continuous variable, discrete+continuous parents (e.g., *Cost*)
- Discrete variable, continuous parents (e.g., *Buy?*)

Chapter 11.1-3 23

Compact conditional distributions

CPT grows exponentially with number of parents

CPT becomes infinite with continuous-valued parent or child

Solution: canonical distributions that are defined compactly

Deterministic nodes are the simplest case:
 $X = f(\text{Parents}(X))$ for some function f

E.g.: Boolean functions

NorthAmerican \Leftrightarrow *Canadian* \vee *US* \vee *Mexican*


E.g.: numerical relationships among continuous variables

$\frac{\partial \text{level}}{\partial t} = \text{inflow} + \text{precipitation} - \text{outflow} - \text{evaporation}$


Chapter 11.1-3 21

Continuous variables

$$\text{Gaussian density } P(x) = \frac{1}{\sqrt{2\pi}\sigma} e^{-\frac{(x-\mu)^2}{2\sigma^2}}$$



Uniform density $P(X=x) = U[18,26](x) = \text{uniform density between 18 and 26}$



Chapter 11.1-3 24

Continuous child variables

Need one **conditional density function** for child variable given continuous parents, for each possible assignment to discrete parents

Most common is the **linear Gaussian model**, e.g.,:

$$\begin{aligned} P(\text{Cost} = c | \text{Harvest} = h, \text{Subsidy} = \text{true}) \\ &= N(a_h h + b_i, \sigma_i)(c) \\ &= \frac{1}{\sigma_i \sqrt{2\pi}} \exp\left(-\frac{1}{2} \left(\frac{c - (a_h h + b_i)}{\sigma_i}\right)^2\right) \end{aligned}$$


Mean *Cost* varies linearly with *Harvest*, variance is fixed

Linear variation is unreasonable over the full range but works OK if the **likely** range of *Harvest* is narrow

Chapter 11.1-3 25


Why the probit?

1. It's sort of the right shape
2. Can view as hard threshold whose location is subject to noise



Chapter 11.1-3 28

Continuous child variables



All-continuous network with LG distributions


⇒ full joint distribution is a multivariate Gaussian

Discrete+continuous LG network is a **conditional Gaussian network** i.e., a multivariate Gaussian over all continuous variables for each combination of discrete variable values

Chapter 11.1-3 26

Discrete variable w/ continuous parents

Probability of *Buys?* given *Cost* should be a "soft" threshold:



Probit distribution uses **integral of Gaussian**:

$$\begin{aligned} \Phi(x) &= \int_{-\infty}^x N(0, 1)(t) dt \\ P(\text{Buys?} = \text{true} | \text{Cost} = c) &= \Phi((-c + \mu)/\sigma) \end{aligned}$$


Chapter 11.1-3 27

Discrete variable contd.

Sigmoid (or logit) distribution also used in neural networks:

$$P(\text{Buys?} = \text{true} | \text{Cost} = c) = \frac{1}{1 + \exp\left(\frac{-2-c\mu}{\sigma}\right)}$$

Sigmoid has similar shape to probit but much longer tails:



Chapter 11.1-3 29

Summary

Bayes nets provide a natural representation for (causally induced) conditional independence

Topology + CPTs = compact representation of joint distribution

Generally easy for (non)experts to construct

Canonical distributions (e.g., noisy-OR) = compact representation of CPTs

Continuous variables ⇒ parameterized distributions (e.g., linear Gaussian)

Chapter 11.1-3 30