

# Graphical Models - Part I

Oliver Schulte - CMPT 726

Bishop PRML Ch. 8, some slides from Russell and Norvig  
AIMA2e

# Outline

Probabilistic Models

Bayesian Networks

Markov Random Fields

Inference

# Outline

Probabilistic Models

Bayesian Networks

Markov Random Fields

Inference

# Probabilistic Models

- We now turn our focus to probabilistic models for pattern recognition
  - Probabilities express beliefs about uncertain events, useful for decision making, combining sources of information
- Key quantity in probabilistic reasoning is the **joint distribution**

$$p(x_1, x_2, \dots, x_K)$$

where  $x_1$  to  $x_K$  are all variables in model

- Address two problems
  - **Inference**: answering queries given the joint distribution
  - **Learning**: deciding what the joint distribution is (involves inference)
- **All inference and learning problems involve manipulations of the joint distribution**

## Reminder - Three Tricks

- Bayes' rule:

$$p(Y|X) = \frac{p(X|Y)p(Y)}{p(X)} = \alpha p(X|Y)p(Y)$$

- Marginalization:

$$p(X) = \sum_y p(X, Y = y) \text{ or } p(X) = \int p(X, Y = y) dy$$

- Product rule:

$$p(X, Y) = p(X)p(Y|X)$$

- All 3 work with extra conditioning, e.g.:

$$p(X|Z) = \sum_y p(X, Y = y|Z)$$

$$p(Y|X, Z) = \alpha p(X|Y, Z)p(Y|Z)$$

# Joint Distribution

	<i>toothache</i>		$\neg$ <i>toothache</i>	
	<i>catch</i>	$\neg$ <i>catch</i>	<i>catch</i>	$\neg$ <i>catch</i>
<i>cavity</i>	<b>.108</b>	<b>.012</b>	<b>.072</b>	<b>.008</b>
$\neg$ <i>cavity</i>	<b>.016</b>	<b>.064</b>	<b>.144</b>	<b>.576</b>

- Consider model with 3 boolean random variables: *cavity*, *catch*, *toothache*
- Can answer query such as

$$p(\neg \text{cavity} | \text{toothache})$$

# Joint Distribution

	<i>toothache</i>		$\neg$ <i>toothache</i>	
	<i>catch</i>	$\neg$ <i>catch</i>	<i>catch</i>	$\neg$ <i>catch</i>
<i>cavity</i>	<b>.108</b>	<b>.012</b>	<b>.072</b>	<b>.008</b>
$\neg$ <i>cavity</i>	<b>.016</b>	<b>.064</b>	<b>.144</b>	<b>.576</b>

- Consider model with 3 boolean random variables: *cavity*, *catch*, *toothache*
- Can answer query such as

$$p(\neg \text{cavity} | \text{toothache})$$

# Joint Distribution

	<i>toothache</i>		$\neg$ <i>toothache</i>	
	<i>catch</i>	$\neg$ <i>catch</i>	<i>catch</i>	$\neg$ <i>catch</i>
<i>cavity</i>	<b>.108</b>	<b>.012</b>	<b>.072</b>	<b>.008</b>
$\neg$ <i>cavity</i>	<b>.016</b>	<b>.064</b>	<b>.144</b>	<b>.576</b>

- Consider model with 3 boolean random variables: *cavity*, *catch*, *toothache*
- Can answer query such as

$$p(\neg\text{cavity}|\text{toothache}) = \frac{p(\neg\text{cavity}, \text{toothache})}{p(\text{toothache})}$$

$$p(\neg\text{cavity}|\text{toothache}) = \frac{0.016 + 0.064}{0.108 + 0.012 + 0.016 + 0.064} = 0.4$$



# Joint Distribution

	<i>toothache</i>		$\neg$ <i>toothache</i>	
	<i>catch</i>	$\neg$ <i>catch</i>	<i>catch</i>	$\neg$ <i>catch</i>
<i>cavity</i>	<b>.108</b>	<b>.012</b>	<b>.072</b>	<b>.008</b>
$\neg$ <i>cavity</i>	<b>.016</b>	<b>.064</b>	<b>.144</b>	<b>.576</b>

- Consider model with 3 boolean random variables: *cavity*, *catch*, *toothache*
- Can answer query such as

$$p(\neg\text{cavity}|\text{toothache}) = \frac{p(\neg\text{cavity}, \text{toothache})}{p(\text{toothache})}$$

$$p(\neg\text{cavity}|\text{toothache}) = \frac{0.016 + 0.064}{0.108 + 0.012 + 0.016 + 0.064} = 0.4$$

# Joint Distribution


- In general, to answer a query on random variables  $\mathbf{Q} = Q_1, \dots, Q_N$  given evidence  $\mathbf{E} = \mathbf{e}$ ,  $\mathbf{E} = E_1, \dots, E_M$ ,  $\mathbf{e} = e_1, \dots, e_M$ :

$$\begin{aligned} p(\mathbf{Q}|\mathbf{E} = \mathbf{e}) &= \frac{p(\mathbf{Q}, \mathbf{E} = \mathbf{e})}{p(\mathbf{E} = \mathbf{e})} \\ &= \frac{\sum_{\mathbf{h}} p(\mathbf{Q}, \mathbf{E} = \mathbf{e}, \mathbf{H} = \mathbf{h})}{\sum_{\mathbf{q}, \mathbf{h}} p(\mathbf{Q} = \mathbf{q}, \mathbf{E} = \mathbf{e}, \mathbf{H} = \mathbf{h})} \end{aligned}$$

# Problems

- The joint distribution is large
  - e. g. with  $K$  boolean random variables,  $2^K$  entries
- Inference is slow, previous summations take  $O(2^K)$  time
- Learning is difficult, data for  $2^K$  parameters
- Analogous problems for continuous random variables

## Reminder - Independence



- $A$  and  $B$  are **independent** iff
 
$$p(A|B) = p(A) \quad \text{or} \quad p(B|A) = p(B) \quad \text{or} \quad p(A, B) = p(A)p(B)$$
- $p(\textit{Toothache}, \textit{Catch}, \textit{Cavity}, \textit{Weather}) = p(\textit{Toothache}, \textit{Catch}, \textit{Cavity})p(\textit{Weather})$ 
  - 32 entries reduced to 12 (*Weather* takes one of 4 values)
- Absolute independence powerful but rare
- Dentistry is a large field with hundreds of variables, none of which are independent. What to do?

## Reminder - Conditional Independence

- $p(\textit{Toothache}, \textit{Cavity}, \textit{Catch})$  has  $2^3 - 1 = 7$  independent entries
- If I have a cavity, the probability that the probe catches in it doesn't depend on whether I have a toothache:
  - (1)  $P(\textit{catch}|\textit{toothache}, \textit{cavity}) = P(\textit{catch}|\textit{cavity})$
- The same independence holds if I haven't got a cavity:
  - (2)  $P(\textit{catch}|\textit{toothache}, \neg\textit{cavity}) = P(\textit{catch}|\neg\textit{cavity})$
- *Catch* is **conditionally independent** of *Toothache* given *Cavity*:  $p(\textit{Catch}|\textit{Toothache}, \textit{Cavity}) = p(\textit{Catch}|\textit{Cavity})$
- Equivalent statements:
  - $p(\textit{Toothache}|\textit{Catch}, \textit{Cavity}) = p(\textit{Toothache}|\textit{Cavity})$
  - $p(\textit{Toothache}, \textit{Catch}|\textit{Cavity}) = p(\textit{Toothache}|\textit{Cavity})p(\textit{Catch}|\textit{Cavity})$
  - $\textit{Toothache} \perp\!\!\!\perp \textit{Catch}|\textit{Cavity}$

## Conditional Independence contd.

- Write out full joint distribution using chain rule:

$$p(\textit{Toothache}, \textit{Catch}, \textit{Cavity})$$

$$= p(\textit{Toothache} | \textit{Catch}, \textit{Cavity})p(\textit{Catch}, \textit{Cavity})$$

$$= p(\textit{Toothache} | \textit{Catch}, \textit{Cavity})p(\textit{Catch} | \textit{Cavity})p(\textit{Cavity})$$

$$= p(\textit{Toothache} | \textit{Cavity})p(\textit{Catch} | \textit{Cavity})p(\textit{Cavity})$$

2 + 2 + 1 = 5 independent numbers

- In many cases, the use of conditional independence greatly reduces the size of the representation of the joint distribution

# Graphical Models

- Graphical Models provide a visual depiction of probabilistic model
- Conditional independence assumptions can be seen in graph
- Inference and learning algorithms can be expressed in terms of graph operations
- We will look at 3 types of graph (can be combined)
  - Directed graphs: [Bayesian networks](#)
  - Undirected graphs: [Markov Random Fields](#)
  - [Factor graphs](#)

# Outline

Probabilistic Models

**Bayesian Networks**

Markov Random Fields

Inference




# Bayesian Networks

- A simple, graphical notation for conditional independence assertions and hence for compact specification of full joint distributions
- Syntax:
  - a set of nodes, one per variable
  - a directed, acyclic graph (link  $\approx$  “directly influences”)
  - a conditional distribution for each node given its parents:

$$p(X_i | pa(X_i))$$

- In the simplest case, conditional distribution represented as a **conditional probability table** (CPT) giving the distribution over  $X_i$  for each combination of parent values

## Example




- Topology of network encodes conditional independence assertions:
  - *Weather* is independent of the other variables
  - *Toothache* and *Catch* are conditionally independent given *Cavity*

## Example


- I'm at work, neighbor John calls to say my alarm is ringing, but neighbor Mary doesn't call. Sometimes it's set off by minor earthquakes. Is there a burglar?
- Variables: *Burglar*, *Earthquake*, *Alarm*, *JohnCalls*, *MaryCalls*
- Network topology reflects causal knowledge:
  - A burglar can set the alarm off
  - An earthquake can set the alarm off
  - The alarm can cause Mary to call
  - The alarm can cause John to call
- (Causal models and conditional independence seem hardwired for humans!)

## Example contd.




# Compactness

- A CPT for Boolean  $X_i$  with  $k$  Boolean parents has  $2^k$  rows for the combinations of parent values
- Each row requires one number  $p$  for  $X_i = true$  (the number for  $X_i = false$  is just  $1 - p$ )
- If each variable has no more than  $k$  parents, the complete network requires  $O(n \cdot 2^k)$  numbers
- i.e., grows linearly with  $n$ , vs.  $O(2^n)$  for the full joint distribution
- For burglary net, ?? numbers



# Compactness

- A CPT for Boolean  $X_i$  with  $k$  Boolean parents has  $2^k$  rows for the combinations of parent values
- Each row requires one number  $p$  for  $X_i = \text{true}$  (the number for  $X_i = \text{false}$  is just  $1 - p$ )
- If each variable has no more than  $k$  parents, the complete network requires  $O(n \cdot 2^k)$  numbers
- i.e., grows linearly with  $n$ , vs.  $O(2^n)$  for the full joint distribution
- For burglary net, ?? numbers
  - $1 + 1 + 4 + 2 + 2 = 10$  numbers (vs.  $2^5 - 1 = 31$ )




# Global Semantics

- **Global semantics** defines the full joint distribution as the product of the local conditional distributions:

$$P(x_1, \dots, x_n) = \prod_{i=1}^n P(x_i | pa(X_i))$$

e.g.,  $P(j \wedge m \wedge a \wedge \neg b \wedge \neg e) =$




# Global Semantics

- **Global semantics** defines the full joint distribution as the product of the local conditional distributions:

$$P(x_1, \dots, x_n) = \prod_{i=1}^n P(x_i | pa(X_i))$$


e.g.,  $P(j \wedge m \wedge a \wedge \neg b \wedge \neg e) =$

$$\begin{aligned} & P(j|a)P(m|a)P(a|\neg b, \neg e)P(\neg b)P(\neg e) \\ = & 0.9 \times 0.7 \times 0.001 \times 0.999 \times 0.998 \\ \approx & 0.00063 \end{aligned}$$





# Example - Car Insurance




<http://aispace.org/bayes>

# Specifying Distributions - Discrete Variables

- Earlier we saw the use of **conditional probability tables** (CPT) for specifying a distribution over discrete random variables with discrete-valued parents
- For a variable with no parents, with  $K$  possible states:


$$p(\mathbf{x}|\boldsymbol{\mu}) = \prod_{k=1}^K \mu_k^{x_k}$$

- e.g.  $p(B) = 0.001^{B_1} 0.999^{B_2}$ ,  
1-of- $K$  representation



# Specifying Distributions - Discrete Variables cont.

- With two variables  $x_1, x_2$  can have two cases



- Dependent

$$\begin{aligned}
 p(x_1, x_2 | \mu) &= p(x_1 | \mu) p(x_2 | x_1, \mu) \\
 &= \left( \prod_{k=1}^K \mu_{k1}^{x_{1k}} \right) \left( \prod_{k=1}^K \prod_{j=1}^K \mu_{kj2}^{x_{1k} x_{2j}} \right)
 \end{aligned}$$

- $K^2 - 1$  free parameters in  $\mu$

- Independent

$$\begin{aligned}
 p(x_1, x_2 | \mu) &= p(x_1 | \mu) p(x_2 | \mu) \\
 &= \left( \prod_{k=1}^K \mu_{k1}^{x_{1k}} \right) \left( \prod_{k=1}^K \mu_{k2}^{x_{2k}} \right)
 \end{aligned}$$

- $2(K - 1)$  free parameters in  $\mu$

# Chains of Nodes




- With  $M$  nodes, could form a chain as shown above
- Number of parameters is:

$$\underbrace{(K - 1)}_{x_1} + (M - 1) \underbrace{K(K - 1)}_{\text{others}}$$

- Compare to:
  - $K^M - 1$  for fully connected graph
  - $M(K - 1)$  for graph with no edges (all independent)

## Sharing Parameters



- Another way to reduce number of parameters is **sharing** parameters (a. k. a. **tying** of parameters)
- Lower graph reuses same  $\mu$  for nodes 2- $M$ 
  - $\mu$  is a random variable in this network, could also be deterministic
- $(K - 1) + K(K - 1)$  parameters

# Specifying Distributions - Continuous Variables


- One common type of conditional distribution for continuous variables is the **linear-Gaussian**

$$p(x_i | pa_i) = \mathcal{N} \left( x_i; \sum_{j \in pa_i} w_{ij} x_j + b_i, v_i \right)$$

- e.g. With one parent *Harvest*:

$$p(c|h) = \mathcal{N}(c; -0.5h + 5, 1)$$

- For harvest  $h$ , mean cost is  $-0.5h + 5$ , variance is 1



## Linear Gaussian

- Interesting fact: if all nodes in a Bayesian Network are linear Gaussian, joint distribution is a multivariate Gaussian.
- Converse is true as well, see Ch.2.3.

$$p(x_i|pa_i) = \mathcal{N} \left( x_i; \sum_{j \in pa_i} w_{ij}x_j + b_i, v_i \right)$$

$$p(x_1, \dots, x_N) = \prod_{i=1}^N \mathcal{N} \left( x_i; \sum_{j \in pa_i} w_{ij}x_j + b_i, v_i \right)$$


- Each factor looks like  $\exp(-\frac{1}{2}(x_i - (\mathbf{w}_i^T \mathbf{x}_{pa_i}) + b_i)^2 / v_i)$ , this product will be another quadratic form of the components of  $\mathbf{x}$ .
- With no links in graph, end up with diagonal covariance matrix
- With fully connected graph, end up with full covariance matrix

# Conditional Independence in Bayesian Networks

- Recall again that  $a$  and  $b$  are conditionally independent given  $c$  ( $a \perp\!\!\!\perp b|c$ ) if
  - $p(a|b, c) = p(a|c)$  or equivalently
  - $p(a, b|c) = p(a|c)p(b|c)$
- Before we stated that links in a graph are  $\approx$  “direct influences”
- We now develop a correct notion of links, in terms of the conditional independences they represent
  - This will be useful for general-purpose inference methods
  - It provides a fast solution to the *relevance problem*: determine whether  $X$  is relevant to  $Y$  given knowledge of  $Z$ .



# A Tale of Three Graphs - Part 1




- The graph above means

$$\begin{aligned} p(a, b, c) &= p(a|c)p(b|c)p(c) \\ p(a, b) &= \sum_c p(a|c)p(b|c)p(c) \\ &\neq p(a)p(b) \text{ in general} \end{aligned}$$

- So  $a$  and  $b$  not independent

# A Tale of Three Graphs - Part 1




- However, conditioned on  $c$

$$p(a, b|c) = \frac{p(a, b, c)}{p(c)} = \frac{p(a|c)p(b|c)p(c)}{p(c)} = p(a|c)p(b|c)$$

- So  $a \perp\!\!\!\perp b|c$

# A Tale of Three Graphs - Part 1



- Note the **path** from  $a$  to  $b$  in the graph
  - When  $c$  is not observed, path is open,  $a$  and  $b$  not independent
  - When  $c$  is observed, path is blocked,  $a$  and  $b$  independent
- In this case  $c$  is **tail-to-tail** with respect to this path

## A Tale of Three Graphs - Part 2




- The graph above means

$$p(a, b, c) = p(a)p(b|c)p(c|a)$$

- Again  $a$  and  $b$  not independent

## A Tale of Three Graphs - Part 2



- However, conditioned on  $c$

$$\begin{aligned}
 p(a, b|c) &= \frac{p(a, b, c)}{p(c)} = \frac{p(a)p(b|c)}{p(c)}p(c|a) \\
 &= \frac{p(a)p(b|c)}{p(c)} \underbrace{\frac{p(a|c)p(c)}{p(a)}}_{\text{Bayes' Rule}} \\
 &= p(a|c)p(b|c)
 \end{aligned}$$


- So  $a \perp\!\!\!\perp b|c$

## A Tale of Three Graphs - Part 2



- As before, the **path** from  $a$  to  $b$  in the graph
  - When  $c$  is not observed, path is open,  $a$  and  $b$  not independent
  - When  $c$  is observed, path is blocked,  $a$  and  $b$  independent
- In this case  $c$  is **head-to-tail** with respect to this path

## A Tale of Three Graphs - Part 3




- The graph above means

$$\begin{aligned} p(a, b, c) &= p(a)p(b)p(c|a, b) \\ p(a, b) &= \sum_c p(a)p(b)p(c|a, b) \\ &= p(a)p(b) \end{aligned}$$

- This time  $a$  and  $b$  are independent

## A Tale of Three Graphs - Part 3




- However, conditioned on  $c$

$$\begin{aligned} p(a, b|c) &= \frac{p(a, b, c)}{p(c)} = \frac{p(a)p(b)p(c|a, b)}{p(c)} \\ &\neq p(a|c)p(b|c) \text{ in general} \end{aligned}$$

- So  $a$  is dependent on  $b$  given  $c$




## A Tale of Three Graphs - Part 3



- The behaviour here is different
  - When  $c$  is not observed, path is blocked,  $a$  and  $b$  independent
  - When  $c$  is observed, path is unblocked,  $a$  and  $b$  not independent
- In this case  $c$  is **head-to-head** with respect to this path
- Situation is in fact more complex, path is unblocked if any **descendent** of  $c$  is observed

## Part 3 - Intuition




- Binary random variables  $B$  (battery charged),  $F$  (fuel tank full),  $G$  (fuel gauge reads full)
- $B$  and  $F$  independent
- But if we observe  $G = 0$  (false) things change
  - e.g.  $p(F = 0|G = 0, B = 0)$  could be less than  $p(F = 0|G = 0)$ , as  $B = 0$  **explains away** the fact that the gauge reads empty
  - Recall that  $p(F|G, B) = p(F|G)$  is another  $F \perp\!\!\!\perp B|G$

# D-separation


- A general statement of conditional independence
- For sets of nodes  $A$ ,  $B$ ,  $C$ , check all paths from  $A$  to  $B$  in graph
- If all paths are **blocked**, then  $A \perp\!\!\!\perp B \mid C$
- Path is blocked if:
  - Arrows meet **head-to-tail** or **tail-to-tail** at a node in  $C$
  - Arrows meet **head-to-head** at a node, and neither node nor any descendent is in  $C$

# Naive Bayes



- Commonly used **naive Bayes** classification model
- Class label  $z$ , features  $x_1, \dots, x_D$
- Model assumes features independent given class label
  - **Tail-to-tail** at  $z$ , blocks path between features

## Markov Blanket



- What is the minimal set of nodes which makes a node  $x_i$  conditionally independent from the rest of the graph?
  - $x_i$ 's parents, children, and children's parents (co-parents)
- Define this set  $MB$ , and consider:

$$\begin{aligned}
 p(x_i | x_{\{j \neq i\}}) &= \frac{p(x_1, \dots, x_D)}{\int p(x_1, \dots, x_D) dx_i} \\
 &= \frac{\prod_k p(x_k | pa_k)}{\int \prod_k p(x_k | pa_k) dx_i}
 \end{aligned}$$

- All factors other than those for which  $x_i$  is  $x_k$  or in  $pa_k$  cancel

# Learning Parameters

- When all random variables are observed in training data, relatively straight-forward
  - Distribution factors, all factors observed
  - e.g. Maximum likelihood used to set parameters of each distribution  $p(x_i|pa_i)$  separately
- When some random variables not observed, it's tricky
  - This is a common case
  - Expectation-maximization (later) is a method for this

# Outline

Probabilistic Models

Bayesian Networks

**Markov Random Fields**

Inference

# Outline

Probabilistic Models

Bayesian Networks

Markov Random Fields

**Inference**